

Mixed regularization methods for the Cauchy problems of the Helmholtz equation

حجت اله شکری کاوه^۱

^۱ کارشناسی ارشد، گروه ریاضی، دانشکده ریاضی و علوم کامپیوتر، دانشگاه صنعتی امیرکبیر، تهران، ایران

چکیده

In this paper, two mixed filter functions are applied for the regularization of the Helmholtz equation. Bound of error is presented under a priori bound respect to the exact solution. At last, the numerical experiment shows the efficiency and effectiveness of these methods.

Keywords: Ill-posed, Helmholtz equation, Regularization method, Cauchy problem, mixed filter function

Introduction

It is well known that inverse problems of elliptic equations are ill-posed in the sense that small noise in input data can introduce arbitrarily large error in the solution [۲۲]. For continuous dependence of the solution on the Cauchy data for satisfying Third Hadamard well-posedness condition, additional conditions are needed. These conditions are bounds for the norm of the solution and noise.

Class of Regularization methods includes a lot of various methods for minimization of noise effects. The classical methods such as, Tikhonov method, truncate singular value decomposition (TSVD), L-curve method and iterative methods can be found in [۷-۱۱].

In the literature inverse problem of the Cauchy problem of Helmholtz equation solved by some numerical methods such as separation of variable, spectral methods, boundary element method (BEM) and solution is severely ill-posed in large scale problems. Almost, the solution is sensitive respect to noise norm. Regularization methods are approached to decrease this affectability.

To vanish ill-posedness of this problem different regularization methods applied by Xiao-Li Feng et al [۳], Hao Cheng et al [۴], H.H. Qin and T. Wei [۵], Xiangtuan Xionget al [۶], Haihua Qin and Jingmei Lu [۲۵]. In classical methods, accuracy is low and for new iterative methods the computational cost is high.

In this paper, two filter functions are utilized from mixing (TSVD), exponential functions and Tikhonov filter. To decrease the truncation error in numerical implementation of Tikhonov method in unbounded problems, problems with a large variance in a spectrum similar to Cauchy problems of Helmholtz and Laplace equations, we can apply TSVD-exponential filtering by using of TSVD method in one part of the spectrum and exponential filtering in another part.

۱. Mathematical problem

We consider Cauchy problem for the Helmholtz equation with Cauchy conditions:

$$\begin{aligned} \Delta w(x, y) + k^2 w(x, y) &= f, & 0 < x < \pi; 0 < y < 1 & \quad (1.1) \\ w(x, 0) &= \varphi(x), & 0 \leq x \leq \pi & \\ w_y(x, 0) &= \psi(x), & 0 \leq x \leq \pi & \\ w(0, y) = w(\pi, y) &= 0, & 0 \leq y \leq 1 & \end{aligned}$$

According to the linearity of the problem (1.1), we divide it into the following two ill-posed problems:

$$\begin{aligned} \Delta u(x, y) + k^2 u(x, y) &= f, & 0 < x < \pi; 0 < y & \\ & & & < 1 & \quad (1.2) \end{aligned}$$

$$u(x, 0) = \varphi(x), \quad 0 \leq x \leq \pi$$

$$u_y(x, 0) = 0, \quad 0 \leq x \leq \pi$$

$$u(0, y) = u(\pi, y) = 0, \quad 0 \leq y \leq 1$$

and

$$\Delta v(x, y) + k^2 v(x, y) = 0, \quad 0 < x < \pi; 0 < y < 1 \quad (1.3)$$

$$v(x, 0) = 0, \quad 0 \leq x \leq \pi$$

$$v_y(x, 0) = \psi(x), \quad 0 \leq x \leq \pi$$

$$v(0, y) = v(\pi, y) = 0, \quad 0 \leq y \leq 1$$

Where $w = u + v$ is solution of the problem (1.1). In problem (1.2) the positive real number k is the wave number, the noisy data $\varphi^\delta(x)$ instead of Cauchy data is available which satisfies:

$$\|\varphi_\delta - \varphi\| \leq \delta \quad (1.4)$$

Suppose that the a priori bound is:

$$\max \|u(\cdot, 1)\| \leq E \quad (1.5)$$

Spectral Galerkin Method (SGM) leads to a solution of problem (1.2) as follows [6]:

$$u(x, y) = \sum_{n=1}^{\infty} \langle \sin(nx), \varphi(x) \rangle \sin(nx) \cosh(\sqrt{n^2 - k^2} y) \quad (1.6)$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product in $L^2(0, \pi)$ which is defined by:

$$\langle \sin(nx), \varphi(x) \rangle = \frac{2}{\pi} \int_0^\pi \varphi(t) \sin(nt) dt \quad (1.7)$$

If $n \leq k$, then $\cosh(\sqrt{n^2 - k^2}y) = \cos(\sqrt{k^2 - n^2}y)$ and singular values of the problem is small and if $n \gg k$ the singular values are exponentially and multiple with noisy data. For overcoming to noise propagation the introduce the regularization methods.

۲. Regularization methods

Optimality of the regularization methods theoretically is presented in [۱] and recently application of regularizations methods are: iterative methods [۲۳], L-curve method [۱۶], landweber method [۱۴], TSVD method [۱۸], optimizational method [۲۰], variational method [۲۱].

Several methods for elliptic equations are presented in [۱۷, ۱۹, ۲۴] and for this problem some methods are applied such as regularization with posterior parameter [۲], Tikhonov method [۱۵]. Some advantages of these methods are shown in [۱۲-۱۴].

For ill-posed inverse problem $Ax = b$ regularization method is defined as:

$$R_\alpha \stackrel{\text{def}}{=} g_\alpha(A^*A)A^* \quad (۲.۱)$$

Where A^* is the adjoint operator of A .

Filter functions are class of functions which satisfy:

$$\lim_{\alpha \rightarrow 0} \phi_\alpha(\lambda) = 1 \quad (۲.۲)$$

The most popular regularization methods by filtering are Tikhonov and TSVD methods. In these methods inverse operators are defined by filter functions:

$$g_\alpha(\lambda) = \frac{1}{\lambda} \phi_\alpha(\lambda) \quad (۲.۳)$$

For $\phi_\alpha(\lambda) = 1$ the inverse operator is ill-condition and solution is naive. Selection of α can be helpful to avoid noise propagation.

At first, we will consider TSVD filter function for this problem:

$$\phi_\alpha(\lambda) = \begin{cases} 1 & \lambda \geq \alpha \\ 0 & \lambda < \alpha \end{cases} \quad (۲.۴)$$

and

$$g_\alpha(\lambda) = \begin{cases} \frac{1}{\lambda} & \lambda \geq \alpha \\ 0 & \lambda < \alpha \end{cases} \quad (۲.۵)$$

where $\lambda = 1/\cosh(\sqrt{n^r - k^r}y)$ is the singular value of the inverse problem.

Suppose x^*, x_α, x^δ denote exact solution, regularization solution and noisy solution of the problem, respectively. The truncation error function (TEF) is defined as:

$$\begin{aligned} r_\alpha(\lambda) \\ = 1 - \phi_\alpha(\lambda) \end{aligned} \quad (2.6)$$

The truncation error depends on (TEF):

$$\begin{aligned} x^* - x_\alpha = \\ r_\alpha(A^*A)x^* \end{aligned} \quad (2.7)$$

In Tikhonov method we choose the filter function as:

$$\begin{aligned} \phi_\alpha(\lambda) \\ = \frac{\lambda^r}{\lambda^r + \alpha^r} \end{aligned} \quad (2.8)$$

This filter is the cumulative distribution function (CDF) of the gaussian (normal) random variable and

$$\begin{aligned} g_\alpha(\lambda) = \frac{1}{\lambda} \phi_\alpha(\lambda) \\ = \frac{\lambda}{\lambda^r + \alpha^r} \end{aligned} \quad (2.9)$$

Is the probability distribution function (PDF) of the Gaussian (normal) random variable. Tikhonov regularization is equal to normalization of the singular value. Equivalently, we can use the exponential form of these filters:

$$\phi_\alpha(\lambda) = \left(1 - e^{-\frac{\lambda}{\alpha}}\right) \quad (2.10)$$

$$\begin{aligned} g_\alpha(\lambda) = \frac{1}{\lambda} \phi_\alpha(\lambda) \\ = \frac{1}{\lambda} \left(1 - e^{-\frac{\lambda}{\alpha}}\right) \end{aligned} \quad (2.11)$$

For exponential random variable (CDF) and (PDF):

$$\begin{aligned} \phi_\alpha(\lambda) \\ = e^{-\frac{\alpha}{\lambda}} \end{aligned} \quad (2.12)$$

$$g_{\alpha}(\lambda) = \frac{1}{\lambda} \phi_{\alpha}(\lambda) = \frac{1}{\lambda} e^{-\frac{\alpha}{\lambda}} \quad (2.13)$$

By this approach we can build different filter functions with considering (CDF) and (PDF) of random variables. But, filter structure should be similar to distribution of the singular values. In the Cauchy problem of Helmholtz equation, the singularity is exponential and exponential filters can be helpful.

Method (I): By use of TSVD method for the first part of the spectrum, we can select the truncation parameter N from (γ, ρ) as follows:

$$\lambda = 1 / \cosh(\sqrt{n^{\gamma} - k^{\gamma}} y) \geq \alpha \Rightarrow N = \lceil \sqrt{(\operatorname{arccosh}(\alpha))^{\gamma} + k^{\gamma}} \rceil \quad (2.14)$$

And in the another part the filter function is equal to the exponential function (γ, ρ) . The solution of problem $(1, \gamma)$ has the form as:

$$u(x, y) = \sum_{n=1}^N \langle \sin(nx), \varphi(x) \rangle \sin(nx) \cosh(\sqrt{n^{\gamma} - k^{\gamma}} y) + \sum_{n=N+1}^{\infty} \langle \sin(nx), \varphi(x) \rangle \sin(nx) \cosh(\sqrt{n^{\gamma} - k^{\gamma}} y) (1 - e^{-1/\alpha \cosh(\sqrt{n^{\gamma} - k^{\gamma}} y)}) \quad (2.15)$$

Method (II): From (γ, ρ) and similar approach, we get:

$$u(x, y) = \sum_{n=1}^N \langle \sin(nx), \varphi(x) \rangle \sin(nx) \cosh(\sqrt{n^{\gamma} - k^{\gamma}} y) + \sum_{n=N+1}^{\infty} \langle \sin(nx), \varphi(x) \rangle \sin(nx) \cosh(\sqrt{n^{\gamma} - k^{\gamma}} y) e^{-\alpha \cosh(\sqrt{n^{\gamma} - k^{\gamma}} y)} \quad (2.16)$$

۳. Error bounds

The error bound for regularization method (I) is obtained as follow :

Lemma (۳, ۱): Suppose that u is the exact solution and u^δ is the regularization solution by method (I) for problem (۱, ۲) also will two conditions (۱, ۴) and (۱, ۵) satisfying. If we select $\alpha = (\frac{\delta}{E})^y$, then for fixed $0 < y < 1$ we get the error bound:

$$\begin{aligned} \|u(x, y) - u^\delta(x, y)\| &= \delta^{1-y} E^y \left(1 + \frac{\gamma \delta}{\|\varphi\|} \right) \end{aligned} \quad (۳.۱)$$

Proof: We compute bound of error in two parts of spectrum:

If $n \leq N$, then

$$\begin{aligned} \|u_N(x, y) - u_N^\delta(x, y)\|^\gamma &= \frac{\pi}{\gamma} \sum_{n=1}^{\infty} |\langle \varphi^\delta - \varphi, \sin(nx) \rangle|^\gamma \cosh^\gamma(\sqrt{n^\gamma - k^\gamma} y) \leq \frac{\delta^\gamma}{\alpha^\gamma} \\ \|u_N(x, y) - u_N^\delta(x, y)\| &\leq \frac{\delta}{\alpha} \\ &= \delta^{1-y} E^y \end{aligned} \quad (۳.۲)$$

If $n > N$, then

$$\begin{aligned} \|u(x, y) - u_N^\delta(x, y)\|^\gamma &= \\ \|u(x, y) - u_N^\delta(x, y)\|^\gamma &= \frac{\pi}{\gamma} \sum_{n=1}^{\infty} |\langle \varphi^\delta - \varphi, \sin(nx) \rangle|^\gamma \cosh^\gamma(\sqrt{n^\gamma - k^\gamma} y) \left(e^{-\frac{\gamma}{\alpha \cosh(\sqrt{n^\gamma - k^\gamma} y)}} \right) \leq \\ \sup_{n > N} \frac{\cosh^\gamma(\sqrt{n^\gamma - k^\gamma} y)}{\cosh^\gamma(\sqrt{n^\gamma - k^\gamma})} &\left(e^{-\frac{\gamma}{\alpha \cosh(\sqrt{n^\gamma - k^\gamma} y)}} \right) E^\gamma \times \frac{|\langle \varphi - \varphi^\delta, \sin(nx) \rangle|^\gamma}{|\langle \varphi, \sin(nx) \rangle|^\gamma} \end{aligned}$$

Since in this part of the spectrum $\frac{\cosh(\sqrt{n^\gamma - k^\gamma} y)}{\cosh(\sqrt{n^\gamma - k^\gamma})} \leq \gamma e^{-\sqrt{n^\gamma - k^\gamma} (1-y)}$ and $\alpha = (\frac{\delta}{E})^y$, we get:

$$\begin{aligned} \|u(x, y) - u_N^\delta(x, y)\| &= \\ \sup_{n > N} \gamma e^{-\sqrt{n^\gamma - k^\gamma} (1-y)} &\left(e^{-\frac{\gamma}{\alpha \cosh(\sqrt{n^\gamma - k^\gamma} y)}} \right) E \frac{\delta}{\|\varphi\|} \leq \frac{\gamma \delta^{1-y} E^y}{\|\varphi\|} \end{aligned} \quad (۳.۳)$$

Adding (۳,۲) and (۳,۳) yields (۳,۱).

For regularization method (II) the error bound satisfies in (۳,۱) and for problem (۱,۳) with Cauchy data ψ bound of error is similar. In more general problem (۱,۱) bound of error for both methods is utilize from summation of these bounds.

Lemma (۳,۲): Suppose that w and w^δ is the exact and regularization solution by each of methods (I) and (II) for problem (۱,۱) respectively, and two conditions (۱,۴) and (۱,۵) are satisfied. If we select $\alpha = (\frac{\delta}{E})^y$, then for fixed $0 < y < 1$ we get the error bound:

$$\begin{aligned} \|w(x,y) - w^\delta(x,y)\| \\ = \gamma \delta^{1-y} E^y \left(1 + \frac{\gamma \delta}{\|\varphi\|} \right) \end{aligned} \quad (۳.۴)$$

۴. Numerical tests

In this section, some numerical examples are presented for verifying the validity of the proposed methods. To compare the exact and approximate solutions root mean square relative error is used:

RMSRE

$$= \frac{\sqrt{\frac{1}{M^2} \sum_{i=1}^M \sum_{j=1}^M (u^\delta(x_i, y_j) - u(x_i, y_j))^2}}{\sqrt{\frac{1}{M^2} \sum_{i=1}^M \sum_{j=1}^M (u(x_i, y_j))^2}}$$

where $x_i = \frac{\pi}{M}(i-1)$, $y_j = \frac{1}{M}(j-1)$, $i = 1, \dots, M$, $j = 1, \dots, M$, $M = 20$

Test. ۱) We consider the following Cauchy problem:

$$\Delta u(x,y) + ku(x,y) = 0, \quad 0 < x < \pi; 0 < y < 1 \quad (۴.۱)$$

$$u(x, 0) = x^2(\pi - x) + \sin(\gamma kx) \cosh(\sqrt{\gamma}k), \quad 0 \leq x \leq \pi$$

$$u_y(x, 0) = 0, \quad 0 \leq x \leq \pi$$

$$u(x, y) = u(\pi, y) = 0, \quad 0 \leq y \leq 1$$

We select

$$\varphi(x) = u(x, 0) = \sum_{n=1}^m \langle u(x, 1), \sin(nx) \rangle \sin(nx) \frac{1}{\cosh(\sqrt{n^2 - k^2})}$$

as Cauchy data for problem $(\xi, 1)$, with $m=36$, ε denotes the error level and noisy Cauchy data is

$$\varphi^\delta = \varphi + \varepsilon(x - \pi)(\pi - x)\sin(x).$$

In Fig 1 we show exact solution and the error between the regularized solution (I) and the exact solution. In Fig 2 we show the numerical results at $y = 1, k = 0.5$ for

$$\varepsilon = 10^{-3}, 10^{-2}, 5 \times 10^{-2}$$





Fig 1. Exact solution u for $k=0.5$

$u^\delta - u$ with $\varepsilon = 10^{-1}$

Table ۱. RMSRE with $k=0.5$ for method (I)

RMSRE	۰.۰۰۰۰۲۰۳۴۹	۰.۰۰۰۰۵۹۱۳۹	۰.۰۰۰۵۰	0.0281

Table ۲. RMSRE with $k=0.5$ for method (II)

RMSRE	۰.۰۰۰۰۲۹۴۵۲	۰.۰۰۰۰۶۹۸۳۸	۰.۰۰۰۴۶	0.0287

The relative root mean square error between the regularized solution methods (I), (II) and the exact solution is shown in Tables ۱, ۲.

Test. ۲) For Cauchy problem:

$$\Delta u(x, y) + ku(x, y) = 0, \quad 0 < x < \pi; 0 < y < 1 \quad (4.2)$$

$$u(x, 0) = \varphi(x) = \sin(x), \quad 0 \leq x \leq \pi$$

$$u_y(x, 0) = 0, \quad 0 \leq x \leq \pi$$

$$u(0, y) = u(\pi, y) = 0, \quad 0 \leq y \leq 1$$

the exact solution is:


$$u(x, y) = \cosh(\sqrt{1 - k^2}y) \sin(x) \quad (4.3)$$

Suppose that the noisy version Cauchy data is $\varphi^\delta = \varphi + \varepsilon \text{randn}(\text{size}(\varphi))$ and $M=0.1$.


In Fig ۳ we show exact and regularization solution for methods (I) and (II) for $k=0.3, 0.7$ and $\varepsilon = 10^{-2}$ at $y=0.9$. the errors for $k=0.7$ and $y=0.9, 0.9$ are shown in Table ۳.

Table ۳. $k=۲,۷, \varepsilon = ۰,۱$


RMSRE (I)	۰,۰۱۲۰	۰,۰۰۶۴
RMSRE (II)	۰,۰۱۲۲	۰,۰۰۶۶



(a) $\varepsilon = 10^{-1}$




(b) $\varepsilon = 10^{-1}$




(c) $\varepsilon = 10^{-5}$

Fig ۲. $K=0.9, y=1$



$k=0.7, y=0.9$



$k=2, y=0.9$

Fig ۳. $\varepsilon = 10^{-1}$


Test. ۳) The exact solution of the problem (۱,۱) with $\varphi = \psi = \sin(\sqrt{\nu}kx)$ and $k = \sqrt{\nu}, \frac{1}{\sqrt{\nu}}$ is:

$$u(x, y) = \sin(\sqrt{\nu}kx) \left(\cosh(ky) + \frac{\sinh(ky)}{k} \right)$$


The numerical results for $k = \frac{1}{\sqrt{x}}$, $\varepsilon = 5 \times 10^{-2}$, $y = 0.1, 0.5, 0.8, 1$ are shown in Fig ۴. and the errors between the exact and regularized solutions for $k = \frac{1}{\sqrt{x}}$, $\varepsilon = 5 \times 10^{-2}$ are shown in Table ۴

Table ۴. $k = \frac{1}{\sqrt{x}}$, $\varepsilon = 5 \times 10^{-2}$


<i>RMSRE(I)</i>	۰.۰۴۰۷	۰.۰۳۹۱	۰.۰۳۳۶	۰.۰۲۸۱	0.0173
<i>RMSRE(II)</i>	۰.۰۴۲۹	۰.۰۴۰۰	۰.۰۳۴۸	۰.۰۳۱۱	0.0179




(a)



(b)



(c)



(d)

Fig ۴. $k = \frac{1}{\sqrt{x}}$, $\epsilon = 10^{-4}$ in the (a) $y = 0.1$ (b) $y = 0.4$ (c) $y = 0.1$ (d) $y = 1$

Conclusion

For both methods the errors are the same. For problems with exponential singularity use of TSVD-exponential filters are efficient. For decreasing truncation error use of exponential filters in truncated domain are effectiveness.

References

- [۱]. Tautenhahn, Ulrich. "Optimality for ill-posed problems under general source conditions." *Numerical Functional Analysis and Optimization* ۱۹,۳-۴ (۱۹۹۸): ۳۷۷-۳۹۸.
- [۲]. Fu, Chu-Li, et al. "A a posteriori regularization for the Cauchy problem for the Helmholtz equation with inhomogeneous Neumann data." *Applied Mathematical Modelling* ۳۹,۱۴ (۲۰۱۵): ۴۱۰۳-۴۱۲۰.

- [۳]. Feng, Xiao-Li, Chu-Li Fu, and Hao Cheng. "A regularization method for solving the Cauchy problem for the Helmholtz equation." *Applied Mathematical Modelling* ۳۵,۷ (۲۰۱۱): ۳۳۰۱-۳۳۱۰.
- [۴]. Cheng, Hao, Chu-Li Fu, and Xiao-Li Feng. "An optimal filtering method for the Cauchy problem of the Helmholtz equation." *Applied Mathematics Letters* ۲۴,۶ (۲۰۱۱): ۹۵۸-۹۶۴.
- [۵]. Qin, H. H., and T. Wei. "Two regularization methods for the Cauchy problems of the Helmholtz equation." *Applied mathematical modelling* ۳۴,۴ (۲۰۱۰): ۹۴۷-۹۶۷.
- [۶]. Xiong, Xiangtuan, Xiaochen Zhao, and Junxia Wang. "Spectral Galerkin method and its application to a Cauchy problem of Helmholtz equation." *Numerical Algorithms* ۶۳,۴ (۲۰۱۳): ۶۹۱-۷۱۱.
- [۷]. Murio, Diego A. *The mollification method and the numerical solution of ill-posed problems*. John Wiley & Sons, ۲۰۱۱.
- [۸]. Engl, H. W., M. Hanke, and A. Neubauer. "Regularization of Inverse Problems. Kluwer Academic Publisher, Dordrecht, Boston, London, ۱۹۹۶."
- [۹]. Tikhonov, A. N., and V. Ya Arsenin. "Solutions of Ill-Posed Problems, VH Winston and Sons, Washington, DC, ۱۹۷۷." *Translated from Russian*.
- [۱۰]. Hansen, Per Christian. *Discrete inverse problems: insight and algorithms*. Vol. ۷. Siam, ۲۰۱۰.
- [۱۱]. Vogel, Curtis R. *Computational methods for inverse problems*. Vol. ۲۳. Siam, ۲۰۰۲.
- [۱۲]. Xiong, Xiangtuan, Wanxia Shi, and Xiaoyan Fan. "Two numerical methods for a Cauchy problem for modified Helmholtz equation." *Applied Mathematical Modelling* ۳۵,۱۰ (۲۰۱۱): ۴۹۵۱-۴۹۶۴.
- [۱۳]. Xiong, Xiang-Tuan. "A regularization method for a Cauchy problem of the Helmholtz equation." *Journal of computational and Applied Mathematics* ۲۳۳,۸ (۲۰۱۰): ۱۷۲۳-۱۷۳۲.
- [۱۴]. Marin, L., et al. "BEM solution for the Cauchy problem associated with Helmholtz-type equations by the Landweber method." *Engineering Analysis with Boundary Elements* ۲۸,۹ (۲۰۰۴): ۱۰۲۵-۱۰۳۴.
- [۱۵]. Qin, H. H., and D. W. Wen. "Tikhonov type regularization method for the Cauchy problem of the modified Helmholtz equation." *Applied Mathematics and Computation* ۲۰۳,۲ (۲۰۰۸): ۶۱۷-۶۲۸.
- [۱۶]. Calvetti, D., et al. "Tikhonov regularization and the L-curve for large discrete ill-posed problems." *Journal of computational and applied mathematics* ۱۲۳,۱-۲ (۲۰۰۰): ۴۲۳-۴۴۶.
- [۱۷]. Tuan, Nguyen Huy, and Daniel Lesnic. "A new general filter regularization method for Cauchy problems for elliptic equations with a locally Lipschitz nonlinear source." *Journal of Mathematical Analysis and Applications* ۴۳۴,۲ (۲۰۱۶): ۱۳۷۶-۱۳۹۳.
- [۱۸]. Zhang, Hongwu, and Ting Wei. "A Fourier truncated regularization method for a Cauchy problem of a semi-linear elliptic equation." *Journal of Inverse and Ill-posed Problems* ۲۲,۲ (۲۰۱۴): ۱۴۳-۱۶۸.
- [۱۹]. Nguyen, Huy Tuan, Quoc Viet Tran, and Van Thinh Nguyen. "Some remarks on a modified Helmholtz equation with inhomogeneous source." *Applied Mathematical Modelling* ۳۷,۳ (۲۰۱۳): ۷۹۳-۸۱۴.
- [۲۰]. Kabanikhin, S. I., and A. L. Karchevsky. "Optimization method for solving the Cauchy problem for an elliptic equation." *Journal of Inverse and Ill-Posed Problems* ۳,۱ (۱۹۹۵): ۲۱-۴۶.
- [۲۱]. Hào, Dinh Nho, et al. "A variational method and approximations of a Cauchy problem for elliptic equations." *Journal of Algorithms & Computational Technology* ۴,۱ (۲۰۱۰): ۸۹-۱۱۹.
- [۲۲]. Alessandrini, Giovanni, et al. "The stability for the Cauchy problem for elliptic equations." *Inverse problems* ۲۵,۱۲ (۲۰۰۹): ۱۲۳۰۰۴.
- [۲۳]. Zhang, H. W., and T. Wei. "Two iterative methods for a Cauchy problem of the elliptic equation with variable coefficients in a strip region." *Numerical algorithms* ۶۵,۴ (۲۰۱۴): ۸۷۵-۸۹۲.
- [۲۴]. Kozlov, V. A., V. G. Maz'ya, and A. V. Fomin. "AN ITERATIVE METHOD FOR SOLVING THE CAUCHY PROBLEM FOR ELLIPTIC EQUATIONS." (۱۹۹۱).
- [۲۵]. Qin, Haihua, and Jingmei Lu. "A modified method for a Cauchy problem of the Helmholtz equation." *Bulletin of the Malaysian Mathematical Sciences Society* ۴۰,۴ (۲۰۱۷): ۱۴۹۳-۱۵۲۲.